

ON GLOBAL INDUCTION MECHANISMS IN A μ -CALCULUS WITH EXPLICIT APPROXIMATIONS*

CHRISTOPH SPRENGER¹ AND MADDS DAM²

Abstract. We investigate a Gentzen-style proof system for the first-order μ -calculus based on cyclic proofs, produced by unfolding fixed point formulas and detecting repeated proof goals. Our system uses explicit ordinal variables and approximations to support a simple semantic induction discharge condition which ensures the well-foundedness of inductive reasoning. As the main result of this paper we propose a new syntactic discharge condition based on traces and establish its equivalence with the semantic condition. We give an automata-theoretic reformulation of this condition which is more suitable for practical proofs. For a detailed comparison with previous work we consider two simpler syntactic conditions and show that they are more restrictive than our new condition.

1991 Mathematics Subject Classification. 68Q60, 03F07, 03B35.

1. INTRODUCTION

The first-order μ -calculus [9] provides a useful setting for semi-automatic program verification. It is expressive enough to encode, from the bottom up, a range of program logics (e.g. LTL, CTL, CTL*, Hoare Logic) as well as process calculi and programming languages including their data types and operational semantics. A framework based on this idea is described by Fredlund [6] and has been applied to a substantial part of the concurrent programming language Erlang in the Erlang Verification Tool [1]. A key aspect in the design of such a framework is proof

* This research was carried out when the first author worked at the Swedish Institute of Computer Science (SICS) in Kista, Sweden, supported by the Swiss European Fellowship No. 83EU-065536. In the final phase of writing he worked at INRIA, supported by the ERCIM fellowship No. 2002-10.

¹ INRIA Sophia Antipolis, 2004, route des Lucioles, BP 93, 06902 Sophia Antipolis, France; e-mail: sprenger@sophia.inria.fr

² Dept. of Microelectronics and Information Technology, Royal Institute of Technology, KTH, Forum 105, 164 40 Kista, Sweden; e-mail: mfd@imit.kth.se

search, in particular the handling of fixed point formulas. The standard approach, Park’s fixed point induction rule (*cf.* [7]), is not suitable for proof search in practise. An alternative is to admit cyclic proof structures (*cf.* [2, 4, 8, 14]) and look for sound *induction discharge conditions*, external global criteria that ensure the well-foundedness of the inductive reasoning. In this setting, proof trees are completed into proof graphs by adding back edges (called repeats) from non-axiom leaves to nodes of which they are substitution instances. This type of proof is favourable to proof search as it allows one to delay decisions concerning induction strategies as long as possible.

In this paper, we study induction discharge conditions in the context of a Gentzen-style proof system for the first-order μ -calculus and present a new syntactic condition, which is weaker than previously published ones in the sense that it qualifies more proof structures as valid proofs. Our proof system is a variant of previous systems [4, 6, 11, 12]. In particular, it shares with [4, 6, 11] the technique, first proposed for the modal μ -calculus by Dam and Gurov [4], of introducing explicit approximation ordinal variables and ordering constraints between them into the proof system. Discharge conditions then rely on these ordering constraints. In the presence of a Cut rule, the use of approximation ordinals considerably simplifies earlier treatments (*cf.* [3]). Dam and Gurov proposed a simple semantic discharge condition, which essentially requires that no infinite path in the proof structure can be assigned a coherent infinite sequence of valuations. This condition expresses in a natural way the requirement of well-foundedness of all inductive reasoning, but due to its semantic nature it is not suitable for the purpose of practical proof. We show that it is equivalent to our practically more useful syntactic condition, while previous syntactic conditions [4, 6, 11] turn out to be strictly more restrictive.

Our new condition relies on the notion of *traces*, which are non-increasing chains (w.r.t. the ordering constraints) of ordinal variables associated with a path of a proof structure. They can be seen as a uniform generalisation of the μ - and ν -traces described by Niwiński and Walukiewicz [8] to systems with a Cut rule and explicit approximants. We identify *progress* in a trace with positions where a strict decrease with respect to the constraints occurs. The equivalence with the semantic condition is then established by showing that a trace progressing at infinitely many positions implies well-foundedness on the semantic side and, conversely, the absence of such a trace gives rise to non-wellfoundedness. Based on the observation that every trace can be transformed into a normal trace where progress is made only at repeat nodes, we are able to give a compact automata-theoretic characterisation of our trace-based discharge condition in terms of an inclusion of the languages recognised by two Büchi automata. This formulation may serve as the basis of an implementation in a proof tool such as the Erlang Verification Tool. Being weaker than previously known conditions, the automata-based criterion might be able to detect proofs where the others fail to do so, which is an advantage for semi-automatic proof search.

For a detailed comparison of our new condition with previously published work, we then turn our attention to two simpler discharge conditions. Common to both of these is that they are based on progress and preservation properties of single

ordinal variables at the repeats of the proof structure. The first condition requires that in each strongly connected subgraph of the proof structure there is a repeat progressing on some ordinal variable, while all other repeats preserve that variable. We show how this condition, which is similar to one proposed by Fredlund [6] (and [12], though in a somewhat different setting), corresponds to a simplified, but strictly stronger version of our automata-based condition. Secondly, we restrict our attention to the special case of *simple* proof structures, where repeats loop back to ancestral nodes (*i.e.* they point to a node on the path from the root to the discharged leaf) and introduce a new alternative notion of discharge where repeats are organised in a partial order, called *induction order*. Progress and preservation properties imposed on each repeat then depend on its position in this order. For simple proof structures this condition generalises the one originally proposed by Dam and Gurov [4] and is equivalent to both the previous one as well as to the condition presented by Schöpp [11]. While drawing its inspiration from the latter, it is more local in the sense that it avoids a quantification over strongly connected subgraphs.

The outline of the rest of the paper is as follows. The next section introduces the μ -calculus with explicit approximations. Section 3 presents the proof system, using the semantic induction discharge condition in the basic notion of proof. In Section 4 we first introduce our trace-based discharge condition and establish its equivalence with the semantic one. Based on the notion of normal traces we then propose an automata-theoretic characterisation of this condition. Restricted forms of syntactic discharge are discussed in Section 5 and compared to previous work. Section 6 concludes the paper with a discussion of the results and an outlook on future work.

2. μ -CALCULUS WITH EXPLICIT APPROXIMATIONS

2.1. FIXED POINTS

We first briefly recall some basic facts from fixed point theory. Suppose A is an arbitrary set. Let $\mathbf{2} = \{0, 1\}$ be the two-point lattice and let $\text{Pred}(A) = \mathbf{2}^A$ be the lattice of predicates over A ordered pointwise.

Definition 2.1. Let $\Psi: \text{Pred}(A) \rightarrow \text{Pred}(A)$ be a monotone map on $\text{Pred}(A)$. The *ordinal approximation* $\mu^\alpha \Psi$ of the fixed point $\mu \Psi$ is defined by

$$\begin{aligned} \mu^0 \Psi &= \lambda x. 0 \\ \mu^{\alpha+1} \Psi &= \Psi(\mu^\alpha \Psi) \\ \mu^\gamma \Psi &= \bigvee_{\alpha < \gamma} \mu^\alpha \Psi \quad \text{for a limit ordinal } \gamma \end{aligned}$$

Theorem 2.2. Let $\Psi: \text{Pred}(A) \rightarrow \text{Pred}(A)$ be a monotone map on $\text{Pred}(A)$. Then

- (1) $\mu \Psi = \bigvee_{\alpha} \mu^\alpha \Psi$ is the least fixed point of Ψ (Knaster-Tarski)
- (2) $\mu^\alpha \Psi = \bigvee_{\beta < \alpha} \Psi(\mu^\beta \Psi)$

2.2. SYNTAX

We assume countably infinite sets of *individual* variables $x, y, z, \dots \in V_I$, *predicate* variables $X^n, Y^n, Z^n, \dots \in V_P^n$ of each arity $n \geq 0$, and *ordinal* variables $\iota, \kappa, \lambda, \dots \in V_O$. We write v, v', \dots for variables of any of the aforementioned types. Let t, t', \dots range over the terms of some first-order signature Σ . We write \bar{t} for a vector t_1, \dots, t_n of terms, let $|\bar{t}|$ denote its length n and $\{\bar{t}\}$ the set $\{t_1, \dots, t_n\}$.

Definition 2.3. The syntax of *μ -calculus formulas ϕ and n -ary abstractions Φ^n over signature Σ* is inductively defined by

$$\begin{aligned} \phi & ::= t = t' \mid \neg\phi \mid \phi_1 \vee \phi_2 \mid \exists x.\phi \mid \exists\kappa.\phi \mid \exists\kappa' < \kappa.\phi \mid \Phi^n(\bar{t}) \\ \Phi^n & ::= X^n \mid \mu X^n(\bar{x}).\phi \mid \mu^\kappa X^n(\bar{x}).\phi \end{aligned}$$

with the restriction that $|\bar{x}| = n$ in *fixed point abstractions* $\mu X^n(\bar{x}).\phi$ and *approximation abstractions* $\mu^\kappa X^n(\bar{x}).\phi$, and $|\bar{t}| = n$ in applications $\Phi^n(\bar{t})$. Furthermore, fixed point and approximation abstraction formation are subject to the usual formal monotonicity condition requiring that each occurrence of X^n in ϕ appears in the scope of an even number of negation symbols.

We will henceforth omit the arity annotations from predicate variables and assume that arities match as required by the previous definition. The sets of free variables of formulas and abstractions are defined as expected. In particular, we have

$$\begin{aligned} \text{fv}(\exists\kappa' < \kappa.\phi) & = (\text{fv}(\phi) - \{\kappa'\}) \cup \{\kappa\} \\ \text{fv}(\Phi(\bar{t})) & = \text{fv}(\Phi) \cup \text{fv}(\bar{t}) \\ \text{fv}(\mu X(\bar{x}).\phi) & = \text{fv}(\phi) - \{X, \bar{x}\} \\ \text{fv}(\mu^\kappa X(\bar{x}).\phi) & = (\text{fv}(\phi) - \{X, \bar{x}\}) \cup \{\kappa\} \end{aligned}$$

This is extended to sets of formulas Δ by defining $\text{fv}(\Delta) = \bigcup\{\text{fv}(\phi) \mid \phi \in \Delta\}$. We identify formulas and abstractions that differ only by a renaming of their bound variables. Dual connectives are defined from the primitive ones in the usual way. The greatest fixed point $\nu X(\bar{x}).\phi$ and the greatest fixed point approximation $\nu^\kappa X(\bar{x}).\phi$ are defined by

$$\begin{aligned} \nu X(\bar{x}).\phi & = \neg\mu X(\bar{x}).\neg\phi[\neg X/X] \\ \nu^\kappa X(\bar{x}).\phi & = \neg\mu^\kappa X(\bar{x}).\neg\phi[\neg X/X] \end{aligned}$$

We assume that substitutions σ, σ', \dots map term variables to terms, predicate variables to abstractions of the same arity and ordinal variables to ordinal variables. We write $\phi\sigma$ or $\sigma(\phi)$ to denote the formula obtained from ϕ by substituting each occurrence of a variable v by $\sigma(v)$, renaming bound variables as necessary to avoid capture of free variables.

2.3. SEMANTICS

Let Σ be a first-order signature. A Σ -model $\mathcal{M} = (\mathcal{A}, \rho)$ consists of a Σ -structure \mathcal{A} interpreting the symbols in Σ and an \mathcal{A} -environment ρ interpreting

each variable in its respective domain. We write $|\mathcal{A}|$ for the support set of the structure \mathcal{A} . The semantics interprets a μ -calculus formula ϕ as an element $\|\phi\|_{\mathcal{M}} \in \mathbf{2}$ and a n -ary abstraction Φ as an element $\|\Phi\|_{\mathcal{M}} \in \text{Pred}(|\mathcal{A}|^n)$. We usually drop \mathcal{M} and write $\|\phi\|_{\rho}$ and $\|\Phi\|_{\rho}$ if the structure \mathcal{A} is clear from the context. The semantics $\|t\|_{\rho} \in |\mathcal{A}|$ of a term t is defined as usual.

Definition 2.4. (Semantics) Given a signature Σ and a Σ -model (\mathcal{A}, ρ) the semantics of μ -calculus formulas ϕ and abstractions Φ over Σ is inductively defined by

$$\begin{aligned}
\|t = t'\|_{\rho} &= \text{if } \|t\|_{\rho} = \|t'\|_{\rho} \text{ then } 1 \text{ else } 0 \\
\|\neg\phi\|_{\rho} &= 1 - \|\phi\|_{\rho} \\
\|\phi_1 \vee \phi_2\|_{\rho} &= \max\{\|\phi_1\|_{\rho}, \|\phi_2\|_{\rho}\} \\
\|\exists x.\phi\|_{\rho} &= \bigvee_{a \in |\mathcal{A}|} \|\phi\|_{\rho[a/x]} \\
\|\exists \kappa.\phi\|_{\rho} &= \bigvee_{\beta} \|\phi\|_{\rho[\beta/\kappa]} \\
\|\exists \kappa' < \kappa.\phi\|_{\rho} &= \bigvee_{\beta < \rho(\kappa)} \|\phi\|_{\rho[\beta/\kappa']} \\
\|\Phi(\bar{t})\|_{\rho} &= \|\Phi\|_{\rho}(\|\bar{t}\|_{\rho}) \\
\|X\|_{\rho} &= \rho(X) \\
\|\mu X(\bar{x}).\phi\|_{\rho} &= \mu\Psi \\
\|\mu^{\kappa} X(\bar{x}).\phi\|_{\rho} &= \mu^{\rho(\kappa)}\Psi
\end{aligned}$$

where $\Psi = \lambda P.\lambda \bar{a}.\|\phi\|_{\rho[P/X, \bar{a}/\bar{x}]}$ in the clauses for fixed point and approximation abstractions.

A model $\mathcal{M} = (\mathcal{A}, \rho)$ *satisfies* a formula ϕ , written $\mathcal{M} \models \phi$ if $\|\phi\|_{\rho} = 1$. The formula ϕ is called *valid*, written $\models \phi$, if it is satisfied in all Σ -models. Given a Σ -model $\mathcal{M} = (\mathcal{A}, \rho)$ we extend ρ a posteriori to terms t and formulas ϕ other than variables by defining $\rho(t) = \|t\|_{\rho}$ and $\rho(\phi) = \|\phi\|_{\rho}$. This allows us to compose substitutions σ and environments ρ as in $\rho \circ \sigma$.

3. THE PROOF SYSTEM

The proof system we present in this section is an adaptation to the first-order setting of the one proposed by Schöpp [11] for the modal μ -calculus. The main difference with the original proposal [4] is the addition of ordinal quantification. We would like to stress, however, that all results of this paper remain valid for systems without ordinal quantification as well as for modal variants.

3.1. ORDINAL CONSTRAINTS

Our proof system uses explicit ordinal approximations of fixed point formulas. Constraints between ordinal variables will be recorded in strict partial orders $\mathcal{O} = (|\mathcal{O}|, <_{\mathcal{O}})$, where $|\mathcal{O}|$ is a finite set of ordinal variables from $V_{\mathcal{O}}$ and $<_{\mathcal{O}}$ is a binary, irreflexive and transitive relation on $|\mathcal{O}|$. We refer to \mathcal{O} as a *set of ordinal constraints*. We write $\iota \leq_{\mathcal{O}} \kappa$ if either $\iota <_{\mathcal{O}} \kappa$ or $\kappa \in |\mathcal{O}|$ and $\iota = \kappa$ (syntactic

identity). If \mathcal{O} and \mathcal{O}' are two strict partial orders we write $\mathcal{O} \subseteq \mathcal{O}'$ to mean that $|\mathcal{O}| \subseteq |\mathcal{O}'|$ and $<_{\mathcal{O}} \subseteq <_{\mathcal{O}'}$. In sequents, we will write \mathcal{O}, κ for $(|\mathcal{O}| \cup \{\kappa\}, <_{\mathcal{O}})$ and $\mathcal{O}, \kappa' < \kappa$ for $\mathcal{O}' = (|\mathcal{O}| \cup \{\kappa', \kappa\}, <_{\mathcal{O}'})$, where $<_{\mathcal{O}'}$ is the transitive closure of $<_{\mathcal{O}} \cup \{(\kappa', \kappa)\}$. It is worth noting that the latter notation will only be used in case \mathcal{O}' is indeed a strict partial order. Given a model $\mathcal{M} = (\mathcal{A}, \rho)$ we say that ρ *respects* \mathcal{O} , if $\rho(\iota) < \rho(\kappa)$ whenever $\iota <_{\mathcal{O}} \kappa$.

3.2. SEQUENTS AND PROOF RULES

The *sequents* of the proof system are of the form $\Gamma \vdash_{\mathcal{O}} \Delta$, where Γ and Δ are finite sets of formulas and $\mathcal{O} = (|\mathcal{O}|, <_{\mathcal{O}})$ is a set of ordinal constraints. A sequent is *well-formed* if all ordinal variables occurring free in Γ or Δ are elements of $|\mathcal{O}|$. We restrict our attention to well-formed sequents without further mention. The set of free variables of a sequent is defined by $\text{fv}(\Gamma \vdash_{\mathcal{O}} \Delta) = \text{fv}(\Gamma \cup \Delta) \cup |\mathcal{O}|$. Given a Σ -model $\mathcal{M} = (\mathcal{A}, \rho)$ we say that \mathcal{M} *satisfies* a sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ whenever ρ respects \mathcal{O} and $\mathcal{M} \models \phi$ for all $\phi \in \Gamma$ then $\mathcal{M} \models \psi$ for some $\psi \in \Delta$. A model \mathcal{M} *falsifies* a sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ if \mathcal{M} does not satisfy $\Gamma \vdash_{\mathcal{O}} \Delta$. The sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ is *valid* if it is satisfied in all models and *invalid* otherwise. The purpose of the proof system is to establish the validity of sequents.

The rules of our proof system are displayed in Tables 1 and 2. They are presented in tableau style with the conclusion above the line and the premises below. Table 1 shows standard rules of first-order logic with equality. The fixed point rules of Table 2 are a direct application of Theorem 2.2. Note the asymmetry of the rules $(\mu_1\text{-L})$ and $(\mu_0\text{-R})$. It is not difficult to show, once the notion of proof has been introduced (see Definition 3.5), the derivability of the sequents

- $\Gamma, (\mu X(\bar{x}).\phi)(\bar{t}) \vdash_{\mathcal{O}} \phi[\mu X(\bar{x}).\phi/X, \bar{t}/\bar{x}], \Delta$, and
- $\Gamma, \exists \kappa. (\mu^{\kappa} X(\bar{x}).\phi)(\bar{t}) \vdash_{\mathcal{O}} (\mu X(\bar{x}).\phi)(\bar{t}), \Delta$

which is sufficient to derive the symmetric rules $(\mu_0\text{-L})$ and $(\mu_1\text{-R})$, respectively. However, rule $(\mu_0\text{-R})$ is not derivable using $(\mu_0\text{-L})$, $(\mu_1\text{-L})$ and $(\mu_1\text{-R})$, essentially because $(\mu_0\text{-R})$ is the only proof rule forcing μ -formulas to be interpreted as fixed points. To see this we use a non-standard interpretation of fixed points¹. Interpret $\mu X(\bar{x}).\phi$ as $\mu^{\omega} X(\bar{x}).\phi$, and interpret ordinal variables as ranging over finite ordinals. This interpretation validates all proof rules including $(\mu_0\text{-L})$, $(\mu_1\text{-L})$ and $(\mu_1\text{-R})$, but not $(\mu_0\text{-R})$, which is seen by considering any model structure and fixed point formula with closure ordinal $\omega + 1$.

Rules $(\exists_{\mathcal{O}}\text{-L})$ and $(\exists_{\mathcal{O}}^{\prec}\text{-L})$ both introduce a fresh ordinal variable while the latter rule additionally generates a new ordinal constraint. Their right hand side versions, $(\exists_{\mathcal{O}}\text{-R})$ and $(\exists_{\mathcal{O}}^{\prec}\text{-R})$, respectively require an ordinal variable and a constraint as a witness. Rule (OrdStr), originally proposed by Schöpp [11], allows us to strengthen ordinal constraints in a controlled way. More precisely, we may add new ordinal variables and constraints to the order \mathcal{O} of a sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ as long as no new variable goes below a variable in $|\mathcal{O}|$. This rule is sometimes helpful to find

¹This argument is contributed by one of the anonymous referees.

Structural Rules	
(Id) $\frac{\Gamma, \phi \vdash_{\mathcal{O}} \phi, \Delta}{\cdot}$	(Cut) $\frac{\Gamma \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi \vdash_{\mathcal{O}} \Delta \quad \Gamma \vdash_{\mathcal{O}} \phi, \Delta}$
(W-L) $\frac{\Gamma, \phi \vdash_{\mathcal{O}} \Delta}{\Gamma \vdash_{\mathcal{O}} \Delta}$	(W-R) $\frac{\Gamma \vdash_{\mathcal{O}} \phi, \Delta}{\Gamma \vdash_{\mathcal{O}} \Delta}$
Logical and Equality Rules	
(\neg -L) $\frac{\Gamma, \neg\phi \vdash_{\mathcal{O}} \Delta}{\Gamma \vdash_{\mathcal{O}} \phi, \Delta}$	(\neg -R) $\frac{\Gamma \vdash_{\mathcal{O}} \neg\phi, \Delta}{\Gamma, \phi \vdash_{\mathcal{O}} \Delta}$
(\vee -L) $\frac{\Gamma, \phi_1 \vee \phi_2 \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi_1 \vdash_{\mathcal{O}} \Delta \quad \Gamma, \phi_2 \vdash_{\mathcal{O}} \Delta}$	(\vee -R) $\frac{\Gamma \vdash_{\mathcal{O}} \phi_1 \vee \phi_2, \Delta}{\Gamma \vdash_{\mathcal{O}} \phi_1, \phi_2, \Delta}$
(\exists -L) $\frac{\Gamma, \exists x. \phi \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi \vdash_{\mathcal{O}} \Delta} \quad x \notin \text{fv}(\Gamma \cup \Delta)$	(\exists -R) $\frac{\Gamma \vdash_{\mathcal{O}} \exists x. \phi, \Delta}{\Gamma \vdash_{\mathcal{O}} \phi[t/x], \Delta}$
(= \mathcal{L}) $\frac{\Gamma[t_2/x], t_1 = t_2 \vdash_{\mathcal{O}} \Delta[t_2/x]}{\Gamma[t_1/x] \vdash_{\mathcal{O}} \Delta[t_1/x]}$	(= \mathcal{R}) $\frac{\Gamma \vdash_{\mathcal{O}} t = t, \Delta}{\cdot}$

TABLE 1. Proof Rules of First-Order Logic with Equality

repeats (see Definition 3.1 below) and seems to be required to prove some simple valid sequents such as $\Gamma, \exists \kappa. \phi \vdash_{\mathcal{O}} \exists \kappa. \exists \kappa' < \kappa. \phi[\kappa'/\kappa], \Delta$.

3.3. PRE-PROOFS, RUNS AND PROOFS

A *derivation tree* $\mathcal{D} = (\mathcal{N}, \mathcal{E}, \mathcal{L})$ is a tree $(\mathcal{N}, \mathcal{E})$ with nodes \mathcal{N} and edges $\mathcal{E} \subseteq \mathcal{N} \times \mathcal{N}$ together with a function \mathcal{L} labelling each node of the tree with a sequent in a way that is consistent with the application of the proof rules. We will often write $N(\Gamma \vdash_{\mathcal{O}} \Delta)$ for $\mathcal{L}(N) = \Gamma \vdash_{\mathcal{O}} \Delta$. The proof structures of our system are essentially finite graphs, which are generated from a derivation tree by adding a back edge from each non-axiom leaf to a node of which it is a repetition (up to some substitution). Let us fix an arbitrary derivation tree $\mathcal{D} = (\mathcal{N}, \mathcal{E}, \mathcal{L})$.

Definition 3.1. (Repeat) Let $M(\Gamma \vdash_{\mathcal{O}} \Delta)$ and $N(\Gamma' \vdash_{\mathcal{O}} \Delta')$ be nodes of \mathcal{D} . Then the triple (M, N, σ) is called a *repeat* of \mathcal{D} , if N is a leaf of \mathcal{D} and σ is a substitution such that

- (1) $\phi \in \Gamma$ implies $\sigma(\phi) \in \Gamma'$, and

Fixed Point Rules	
$(\mu_1\text{-L}) \frac{\Gamma, (\mu X(\bar{x}).\phi)(\bar{t}) \vdash_{\mathcal{O}} \Delta}{\Gamma, \exists \kappa. (\mu^\kappa X(\bar{x}).\phi)(\bar{t}) \vdash_{\mathcal{O}} \Delta}$	$(\mu_0\text{-R}) \frac{\Gamma \vdash_{\mathcal{O}} (\mu X(\bar{x}).\phi)(\bar{t}), \Delta}{\Gamma \vdash_{\mathcal{O}} \phi[\mu X(\bar{x}).\phi/X, \bar{t}/\bar{x}], \Delta}$
$(\mu^\kappa\text{-L}) \frac{\Gamma, (\mu^\kappa X(\bar{x}).\phi)(\bar{t}) \vdash_{\mathcal{O}} \Delta}{\Gamma, \exists \kappa' < \kappa. \phi[\mu^{\kappa'} X(\bar{x}).\phi/X, \bar{t}/\bar{x}] \vdash_{\mathcal{O}} \Delta}$	
$(\mu^\kappa\text{-R}) \frac{\Gamma \vdash_{\mathcal{O}} (\mu^\kappa X(\bar{x}).\phi)(\bar{t}), \Delta}{\Gamma \vdash_{\mathcal{O}} \exists \kappa' < \kappa. \phi[\mu^{\kappa'} X(\bar{x}).\phi/X, \bar{t}/\bar{x}], \Delta}$	
Ordinal Rules	
$(\exists_{\mathcal{O}}\text{-L}) \frac{\Gamma, \exists \kappa. \phi \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi \vdash_{\mathcal{O}, \kappa} \Delta} \quad \kappa \notin \mathcal{O} $	$(\exists_{\mathcal{O}}\text{-R}) \frac{\Gamma \vdash_{\mathcal{O}} \exists \kappa. \phi, \Delta}{\Gamma \vdash_{\mathcal{O}} \phi[\iota/\kappa], \Delta} \quad \iota \in \mathcal{O} $
$(\exists_{\mathcal{O}}^<\text{-L}) \frac{\Gamma, \exists \kappa' < \kappa. \phi \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi \vdash_{\mathcal{O}, \kappa' < \kappa} \Delta} \quad \kappa' \notin \mathcal{O} $	$(\exists_{\mathcal{O}}^<\text{-R}) \frac{\Gamma \vdash_{\mathcal{O}} \exists \kappa' < \kappa. \phi, \Delta}{\Gamma \vdash_{\mathcal{O}} \phi[\iota/\kappa'], \Delta} \quad \iota <_{\mathcal{O}} \kappa$
$(\text{OrdStr}) \frac{\Gamma \vdash_{\mathcal{O}} \Delta}{\Gamma \vdash_{\mathcal{O}'} \Delta} \quad \mathcal{O} \subseteq \mathcal{O}' \text{ and } \iota <_{\mathcal{O}'} \kappa, \kappa \in \mathcal{O} \Rightarrow \iota \in \mathcal{O} $	

TABLE 2. Fixed Point and Ordinal Proof Rules

- (2) $\psi \in \Delta$ implies $\sigma(\psi) \in \Delta'$, and
- (3) $\kappa \in |\mathcal{O}|$ implies $\sigma(\kappa) \in |\mathcal{O}'|$, and
- (4) $\iota <_{\mathcal{O}} \kappa$ implies $\sigma(\iota) <_{\mathcal{O}'} \sigma(\kappa)$.

The node N is called *repeat node* and M its *companion node*.

It is worth remarking that we do not require companions to be ancestors of their corresponding repeat nodes in \mathcal{D} .

Definition 3.2. (Pre-Proof) A *pre-proof* $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ for a sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ consists of a derivation tree $\mathcal{D} = (\mathcal{N}, \mathcal{E}, \mathcal{L})$ with root node labelled by $\Gamma \vdash_{\mathcal{O}} \Delta$ and a set of repeats \mathcal{R} for \mathcal{D} such that each non-axiom leaf appears in exactly one repeat of \mathcal{R} . The *pre-proof graph* of \mathcal{P} is defined by $\mathcal{G}(\mathcal{P}) = (\mathcal{N}, \mathcal{E}', \mathcal{L})$, where $\mathcal{E}' = \mathcal{E} \cup \{(N, M) \mid \exists \sigma. (M, N, \sigma) \in \mathcal{R}\}$.

By a *path* of \mathcal{P} we mean a path in $\mathcal{G}(\mathcal{P})$. A path π is called *rooted* if its first node $\pi(0)$ is the root of \mathcal{D} . We write π^i for the i th suffix of π , that is, the path obtained by dropping the first i nodes of π . This yields the empty sequence in

case that i is greater or equal to the length of π . We say that π *traverses a repeat* $R = (M, N, \sigma)$ if $\pi(i) = N$ and $\pi(i + 1) = M$ for some position i .

Example 3.3. (Lexicographic Order) In order to illustrate the preceding definitions, we now present a proof in our system showing that the lexicographic ordering of two well-founded relations is again well-founded. We assume that our signature includes the unary function symbols π_1 and π_2 (to be interpreted as the left and right projections of a pair). In this example, we write t^1 as a shorthand for $\pi_1(t)$ and t^2 for $\pi_2(t)$.

Well-foundedness can be defined in terms of the notion of accessibility of an element x with respect to a binary relation X :

$$\begin{aligned} \text{Acc}(X, x) &= (\mu Z(z). \forall y. \neg X(y, z) \vee Z(y))(x) \\ \text{Wf}(X) &= \forall x. \text{Acc}(X, x) \end{aligned}$$

The least fixed point in the definition of accessibility forces the absence of infinitely descending X -chains from a given element x of the domain. A binary relation X is well-founded if all elements of the domain are accessible with respect to X . The lexicographic ordering of two binary relations X and Y is defined by

$$\text{Lex}(X, Y)(u, w) = X(u^1, w^1) \vee (u^1 = w^1 \wedge Y(u^2, w^2))$$

With these definitions the sequent we would like to prove valid is

$$\Gamma_0 \vdash \text{Wf}(\text{Lex}(X, Y)) \quad (1)$$

where $\Gamma_0 = \text{Wf}(X), \text{Wf}(Y)$. Before presenting a proof of this sequent, we need to introduce some derived rules and abbreviations. The derivation uses the rules (\wedge -L), (\forall_I -L) and (\forall_I -R) for conjunction and universal quantification:

$$\begin{array}{lll} (\wedge\text{-L}) \frac{\Gamma, \phi_1 \wedge \phi_2 \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi_1, \phi_2 \vdash_{\mathcal{O}} \Delta} & (\forall_I\text{-L}) \frac{\Gamma, \forall x. \phi \vdash_{\mathcal{O}} \Delta}{\Gamma, \phi[t/x] \vdash_{\mathcal{O}} \Delta} & (\forall_I\text{-R}) \frac{\Gamma \vdash_{\mathcal{O}} \forall x. \phi, \Delta}{\Gamma \vdash_{\mathcal{O}} \phi, \Delta} C \end{array}$$

The side condition C of rule (\forall_I -R) requires that $x \notin \text{fv}(\Gamma \cup \Delta)$. These rules are easily derived from their duals (\vee -R), (\exists_I -R) and (\exists_I -L), respectively. We also introduce the following abbreviations:

$$\begin{aligned} \text{Acc}^\kappa(X, x) &= (\mu^\kappa Z(z). \forall y. \neg X(y, z) \vee Z(y))(x) \\ A(X, Y, z) &= \text{Acc}(\text{Lex}(X, Y), z) \end{aligned}$$

Figure 1 shows a derivation tree for sequent (1) which is continued in Figures 2 and 3. For brevity, we use a minimalistic notation for ordinal constraints. We write for instance $\Gamma \vdash_{\lambda' < \lambda}^\kappa \Delta$ for the sequent $\Gamma \vdash_{\mathcal{O}} \Delta$, where $\mathcal{O} = (\{\kappa, \lambda, \lambda'\}, \{(\lambda', \lambda)\})$. This should not give rise to any confusion. We use the label (FO) to denote an unspecified series of first-order logic rule applications.

Let us now look at the derivation in some detail. At the root node N_0 we remove the universal quantifier appearing in the definition of Wf on the right

$$\begin{array}{c}
\frac{N_0[\Gamma_0 \vdash \text{Wf}(\text{Lex}(X, Y))]}{N_1[\Gamma_0 \vdash A(X, Y, w)]} (\forall_I\text{-R}) \\
\frac{}{N_2^*[\Gamma_0, \text{Acc}^\kappa(X, w^1) \vdash_\kappa A(X, Y, w)]} (\text{RS1}) \\
\frac{N_3^{**}[\Gamma_0, \text{Acc}^\kappa(X, w^1), \text{Acc}^\lambda(Y, w^2) \vdash_{\kappa, \lambda} A(X, Y, w)]}{N_4[\Gamma_0, \text{Acc}^\kappa(X, w^1), \text{Acc}^\lambda(Y, w^2), \text{Lex}(X, Y)(u, w) \vdash_{\kappa, \lambda} A(X, Y, u)]} (\text{RS1}) \\
\frac{}{N_4[\Gamma_0, \text{Acc}^\kappa(X, w^1), \text{Acc}^\lambda(Y, w^2), \text{Lex}(X, Y)(u, w) \vdash_{\kappa, \lambda} A(X, Y, u)]} (\mu_0\text{-R, FO}) \\
\frac{}{N_4[\Gamma_0, \text{Acc}^\kappa(X, w^1), \text{Acc}^\lambda(Y, w^2), \text{Lex}(X, Y)(u, w) \vdash_{\kappa, \lambda} A(X, Y, u)]} (\text{FO}) \\
\mathcal{D}_L \qquad \qquad \qquad \mathcal{D}_R
\end{array}$$

FIGURE 1. Derivation tree \mathcal{D} for lexicographic order example

$$\begin{array}{c}
\frac{N_5[\Gamma_0, \text{Acc}^\kappa(X, w^1), X(u^1, w^1) \vdash_{\kappa, \lambda} A(X, Y, u)]}{N_6[\Gamma_0, \neg X(u^1, w^1) \vee \text{Acc}^{\kappa'}(X, u^1), X(u^1, w^1) \vdash_{\kappa' < \kappa} A(X, Y, u)]} (\text{RS2}) \\
\frac{N_7^*[\Gamma_0, \text{Acc}^{\kappa'}(X, u^1) \vdash_{\kappa' < \kappa} A(X, Y, u)] \quad N_8[X(u^1, w^1) \vdash_{\kappa' < \kappa} X(u^1, w^1)]}{N_7^*[\Gamma_0, \text{Acc}^{\kappa'}(X, u^1) \vdash_{\kappa' < \kappa} A(X, Y, u)] \quad N_8[X(u^1, w^1) \vdash_{\kappa' < \kappa} X(u^1, w^1)]} (\text{FO})
\end{array}$$

FIGURE 2. Derivation \mathcal{D}_L

$$\begin{array}{c}
\frac{N_9[\Gamma_0, \text{Acc}^\kappa(X, w^1), \text{Acc}^\lambda(Y, w^2), u^1 = w^1, Y(u^2, w^2) \vdash_{\kappa, \lambda} A(X, Y, u)]}{N_{10}[\Gamma_0, \text{Acc}^\kappa(X, u^1), \text{Acc}^\lambda(Y, w^2), Y(u^2, w^2) \vdash_{\kappa, \lambda} A(X, Y, u)]} (= \text{-L}) \\
\frac{N_{11}[\Gamma_0, \text{Acc}^\kappa(X, u^1), \neg Y(u^2, w^2) \vee \text{Acc}^{\lambda'}(Y, u^2), Y(u^2, w^2) \vdash_{\lambda' < \lambda} A(X, Y, u)]}{N_{12}^{**}[\Gamma_0, \text{Acc}^\kappa(X, u^1), \text{Acc}^{\lambda'}(Y, u^2) \vdash_{\lambda' < \lambda} A(X, Y, u)] \quad N_{13}[Y(u^2, w^2) \vdash_{\lambda' < \lambda} Y(u^2, w^2)]} (\text{RS2})
\end{array}$$

FIGURE 3. Derivation \mathcal{D}_R

hand side of the turnstile. Then we apply rule sequence (RS1)=($\forall_I\text{-L}$, $\mu_1\text{-L}$, $\exists_O\text{-L}$) twice to approximate $\text{Acc}(X, w^1)$ to $\text{Acc}^\kappa(X, w^1)$ at node N_1 and $\text{Acc}(Y, w^2)$ to $\text{Acc}^\lambda(Y, w^2)$ at N_2 , introducing the fresh ordinal variables κ and λ , respectively. At node N_3 the formula $A(X, Y, w)$ is unfolded using rule ($\mu_0\text{-R}$) followed by applications of first-order logic rules. Next, at N_4 we apply a series of boolean rules to $\text{Lex}(X, Y)(u, w)$, producing nodes N_5 in Figure 2 and N_9 in Figure 3, corresponding to the two cases in the definition of the lexicographic ordering. From N_5 to N_7 we unfold and instantiate the approximation $\text{Acc}^\kappa(X, w^1)$ using rule sequence (RS2)=($\mu^\kappa\text{-L}$, $\exists_O^\kappa\text{-L}$, $\forall_I\text{-L}$) and first-order logic rules. This yields a new ordinal constraint $\kappa' < \kappa$ and approximation $\text{Acc}^{\kappa'}(X, u^1)$ as well as the axiom node N_7 . In \mathcal{D}_R , after rewriting the equation at node N_{11} , we apply the same sequence of rules as in \mathcal{D}_L to $\text{Acc}^\lambda(Y, w^2)$, resulting in the ordinal constraint $\lambda' < \lambda$ and the approximation $\text{Acc}^{\lambda'}(Y, u^2)$ at node N_{12} and the axiom at N_{13} .

Finally, we extend the derivation tree \mathcal{D} with two repeats L and R (indicated in the figures by $*$ and $**$) as follows:

$$\begin{aligned} L &= (N_2, N_7, \sigma_L) \quad \text{where } \sigma_L = [u/w, \kappa'/\kappa] \\ R &= (N_3, N_{12}, \sigma_R) \quad \text{where } \sigma_R = [u/w, \lambda'/\lambda] \end{aligned}$$

This yields the pre-proof $\mathcal{P} = (\mathcal{D}, \{L, R\})$.

Not every pre-proof is a proof of the validity of its root sequent. The simplest example of a pre-proof that is not a proof is given by the derivation that infers the sequent $\vdash \mu X.X$ from itself using rule $(\mu_0\text{-R})$. This two-node pre-proof clearly uses circular reasoning and should therefore be rejected as a proof.

We now give a simple semantic *induction discharge condition* ensuring that all inductive reasoning embodied in a pre-proof is well-founded.

Definition 3.4. (Run) Let $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ be a pre-proof, \mathcal{A} a Σ -structure and suppose $\Pi = (N_0, \rho_0) \cdots (N_i, \rho_i) \cdots$ is a (finite or infinite) sequence of pairs of nodes of \mathcal{D} and \mathcal{A} -environments. Suppose that $N_i(\Gamma_i \vdash_{\mathcal{O}_i} \Delta_i)$. Then Π is called a *run of the pre-proof* \mathcal{P} if

- (1) N_0 is the root of \mathcal{P} , and
- (2) ρ_i respects \mathcal{O}_i for all i , and
- (3) for all successive pairs (N_i, N_{i+1}) of nodes, either
 - (a) $(N_i, N_{i+1}) \in \mathcal{E}$ and ρ_{i+1} agrees with ρ_i on all free variables common to N_i and N_{i+1} , or
 - (b) $(N_{i+1}, N_i, \sigma) \in \mathcal{R}$ and $\rho_{i+1} = \rho_i \circ \sigma$.

Note that $\pi = N_0 \cdots N_i \cdots$ is a rooted path of \mathcal{P} . We say that the *run* Π *follows the path* π .

Definition 3.5. (Proof) A pre-proof \mathcal{P} for $\Gamma \vdash_{\mathcal{O}} \Delta$ satisfies *discharge condition (rDC)* if all runs of \mathcal{P} are finite, in which case \mathcal{P} is called a *proof for* $\Gamma \vdash_{\mathcal{O}} \Delta$.

Intuitively, the finiteness of runs in a proof rests on the well-foundedness of the underlying interpretation of the ordinal variables and thus prevents unsound circular reasoning. Note that the finiteness of runs is independent of the interpretation of non-ordinal variables. This intuition will be made explicit in Section 4.

Example 3.6. Consider again the pre-proof \mathcal{P} from Example 3.3. Let $\pi_L = N_2 \cdots N_7$ and $\pi_R = N_3 N_4 N_9 \cdots N_{12}$ be the simple cycles corresponding to repeats L and R , respectively. Suppose there is an infinite run r following a path traversing the left loop indefinitely from some point on, that is, a path ending in π_L^ω . By Definition 3.4 the interpretation of κ remains constant along π_L , since κ is free in all of these sequents. Now suppose $r(i) = (N_7, \rho_i)$ and $r(i+1) = (N_2, \rho_{i+1})$. This step traverses the repeat L . In this case Definition 3.4 requires that $\rho_{i+1}(\kappa) = (\rho_i \circ \sigma_L)(\kappa) = \rho_i(\kappa')$. But since ρ_i respects $\kappa' < \kappa$ at N_7 , we have $\rho_i(\kappa') < \rho_i(\kappa)$ and hence $\rho_{i+1}(\kappa) < \rho_i(\kappa)$. Since the run r was assumed to be infinite and traverse L infinitely often, the interpretation of κ must decrease at infinitely many positions. This contradicts the well-foundedness of the ordinals. Hence, such a run does not

exist. A similar argument shows that there is no infinite run following a path ending in π_R^ω , since this implies that the interpretation of λ decreases indefinitely.

By observing that the interpretation of κ remains constant along π_R and when traversing repeat R , we see that there is no infinite run following any path traversing each of the loops π_L and π_R infinitely often. Thus, \mathcal{P} is a proof. We conclude this example by remarking that the syntactic discharge conditions introduced in Sections 4 and 5 provide a much more convenient method to determine whether a pre-proof is a proof.

3.4. SOUNDNESS

Lemma 3.7. (Local soundness) *The proof rules of Tables 1 and 2 are sound. In particular, if there is a Σ -model (\mathcal{A}, ρ) falsifying the conclusion C of a rule then there is an \mathcal{A} -environment ρ' such that (\mathcal{A}, ρ') falsifies some premise P of that rule. Moreover, ρ and ρ' agree on all free variables common to C and P .*

Proof. By inspection of the proof rules. For the fixed point rules the claim follows immediately from Theorem 2.2. The cases of the ordinal rules are straightforward except for rule (OrdStr), which we discuss now. Suppose some model (\mathcal{A}, ρ) falsifies the sequent $\Gamma \vdash_{\mathcal{O}} \Delta$ in the conclusion. The sequent in the premise is $\Gamma \vdash_{\mathcal{O}'} \Delta$ with $\mathcal{O} \subseteq \mathcal{O}'$. Since $<_{\mathcal{O}'}$ is a finite strict partial order, we can define the environment ρ' by well-founded induction on $<_{\mathcal{O}'}$ as follows

$$\rho'(v) = \begin{cases} \max\{\rho'(\iota) \mid \iota <_{\mathcal{O}'} v\} + 1 & \text{if } v \in |\mathcal{O}'| - |\mathcal{O}| \\ \rho(v) & \text{otherwise} \end{cases}$$

It is sufficient to check that ρ' respects \mathcal{O}' and thus falsifies the premise sequent $\Gamma \vdash_{\mathcal{O}'} \Delta$. Consider some constraint $\iota <_{\mathcal{O}'} \kappa$. If $\kappa \in |\mathcal{O}|$ then also $\iota \in |\mathcal{O}|$ by the side condition of the rule. Hence, $\rho'(\iota) < \rho'(\kappa)$ is inherited from \mathcal{O} . If $\kappa \in |\mathcal{O}'| - |\mathcal{O}|$ then we have $\rho'(\iota) < \rho'(\iota) + 1 \leq \rho'(\kappa)$ by the definition of ρ' . \square

Lemma 3.8. *Let (M, N, σ) be a repeat and let (\mathcal{A}, ρ) be a model falsifying $\mathcal{L}(N)$. Then $(\mathcal{A}, \rho \circ \sigma)$ falsifies $\mathcal{L}(M)$.*

Proof. By the definition of a repeat. \square

Theorem 3.9. (Soundness) *If there is a proof for $\Gamma \vdash_{\mathcal{O}} \Delta$, then $\Gamma \vdash_{\mathcal{O}} \Delta$ is valid.*

Proof. Let $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ be a proof for $\Gamma \vdash_{\mathcal{O}} \Delta$ and suppose for a contradiction that some model (\mathcal{A}, ρ_0) falsifies $\Gamma \vdash_{\mathcal{O}} \Delta$. We iteratively construct an infinite run Π . The construction starts with $\Pi_0 = (N_0, \rho_0)$, where N_0 is the root node of \mathcal{P} . Clearly, ρ_0 respects \mathcal{O}_0 since (\mathcal{A}, ρ_0) falsifies $\mathcal{L}(N_0)$. Hence, Π_0 is a finite run.

Assume that we have already constructed the finite run $\Pi_i = (N_0, \rho_0) \cdots (N_i, \rho_i)$ for some $i \geq 0$ such that (\mathcal{A}, ρ_i) is a model falsifying $\mathcal{L}(N_i)$. Note that $\mathcal{L}(N_i)$ can not be an axiom. We show that Π_i can be extended to a run $\Pi_{i+1} = \Pi_i(N_{i+1}, \rho_{i+1})$ such that $(\mathcal{A}, \rho_{i+1})$ is a model falsifying $\mathcal{L}(N_{i+1}) = \Gamma_{i+1} \vdash_{\mathcal{O}_{i+1}} \Delta_{i+1}$ (and hence ρ_{i+1} respects \mathcal{O}_{i+1}). We distinguish two cases. If N_i is a non-leaf node then there

is by Lemma 3.7 a successor node M of N_i and an \mathcal{A} -environment ρ such that (\mathcal{A}, ρ) falsifies $\mathcal{L}(M)$ and ρ agrees with ρ_i on all free variables common to $\mathcal{L}(N_i)$ and $\mathcal{L}(M)$. Then the sequence Π_{i+1} obtained by setting $N_{i+1} = M$ and $\rho_{i+1} = \rho$ is again a run. If N_i is a repeat node then there is a repeat $(M, N_i, \sigma) \in \mathcal{R}$ and we set $N_{i+1} = M$ and $\rho_{i+1} = \rho_i \circ \sigma$. By Lemma 3.8 $(\mathcal{A}, \rho_{i+1})$ falsifies $\mathcal{L}(N_{i+1})$, so Π_{i+1} is a run. The limit Π of the sequence Π_0, Π_1, \dots is an infinite run of \mathcal{P} , contradicting the assumption that \mathcal{P} is a proof. \square

4. SYNTACTIC DISCHARGE CONDITIONS

As condition (rDC) captures the well-foundedness of the reasoning in a pre-proof in a very natural way, it serves as our reference discharge condition. Due to its semantic nature it is, however, hardly usable in practical proofs and we therefore introduce two alternative, purely syntactical, discharge conditions and show that they characterise condition (rDC). For the remainder of this section we fix an arbitrary pre-proof $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ with $\mathcal{D} = (\mathcal{N}, \mathcal{E}, \mathcal{L})$.

4.1. TRACES AND PROGRESS

A trace of \mathcal{P} is a path of \mathcal{P} labelled by ordinal constraints that are linked to yield a non-increasing chain of ordinal dependencies.

Definition 4.1. (Trace) Let $\tau = (N_0, (\kappa_0, \kappa'_0)) \cdots (N_i, (\kappa_i, \kappa'_i)) \cdots$ be a (finite or infinite) sequence of pairs consisting of a node of \mathcal{D} and a pair of ordinal variables of $V_{\mathcal{O}}$. Suppose that $N_i(\Gamma_i \vdash_{\mathcal{O}_i} \Delta_i)$. Then τ is a *trace* of \mathcal{P} if

- (1) $\kappa'_i \leq_{\mathcal{O}_i} \kappa_i$ for all i , and
- (2) for all successive pairs (N_i, N_{i+1}) of nodes, either
 - (a) $(N_i, N_{i+1}) \in \mathcal{E}$ and $\kappa'_i = \kappa_{i+1}$, or
 - (b) $(N_{i+1}, N_i, \sigma) \in \mathcal{R}$ and $\kappa'_i = \sigma(\kappa_{i+1})$.

We say that the trace $\tau = (N_0, (\kappa_0, \kappa'_0)) \cdots (N_i, (\kappa_i, \kappa'_i)) \cdots$ follows the path $\pi = N_0 \cdots N_i \cdots$.

Definition 4.2. (Progress) Let $\tau = (N_0, (\kappa_0, \kappa'_0)) \cdots (N_i, (\kappa_i, \kappa'_i)) \cdots$ and suppose $\mathcal{L}(N_i) = \Gamma_i \vdash_{\mathcal{O}_i} \Delta_i$. Then

- τ *progresses at position i* if $\kappa'_i <_{\mathcal{O}_i} \kappa_i$, and
- τ is *progressive* if there are infinitely many positions where τ progresses, and
- a path π in $\mathcal{G}(\mathcal{P})$ is *progressive* if there is a progressive trace τ following a suffix π^i of π .

Example 4.3. Figure 4 represents the trace

$$\tau = (N_0, (\delta, \varepsilon))(N_1, (\alpha, \beta))(N_2, (\beta, \gamma))(N_3, (\gamma, \gamma))(N_4, (\kappa, \kappa)),$$




FIGURE 4. Example of a trace

where $R_1 = (N_0, N_1, \sigma_1)$ and $R_2 = (N_3, N_4, \sigma_2)$ are repeats with $\sigma_1(\alpha) = \epsilon$ and $\sigma_2(\kappa) = \gamma$. The trace follows the path $N_0N_1N_2N_3N_4$ and progresses at positions 0, 1 and 2.

4.2. THE TRACE-BASED DISCHARGE CONDITION

Definition 4.4. (tDC) A pre-proof \mathcal{P} satisfies *discharge condition (tDC)* if all infinite paths of \mathcal{P} are progressive.

Theorem 4.5. *A pre-proof \mathcal{P} is a proof if and only if it satisfies condition (tDC).*

Proof. It is sufficient to show for any infinite rooted path π in \mathcal{P} that there is no infinite run following π if and only if there is a progressive trace following some suffix of π . Let $\pi = N_0 \cdots N_i \cdots$ be an infinite path in \mathcal{P} and let \mathcal{O}_j denote the partial order appearing in sequent $\mathcal{L}(N_j)$.

“ \Rightarrow ” By contraposition. Suppose that there is no progressive trace following a suffix of π . We will construct an infinite run Π following π . Define the height $h_i(\kappa)$ of ordinal variable κ in the order \mathcal{O}_i inductively as follows:

$$h_i(\kappa) = \begin{cases} 0 & \text{if } \kappa \text{ is minimal in } \mathcal{O}_i \\ \max\{h_i(\iota) \mid \iota <_{\mathcal{O}_i} \kappa\} + 1 & \text{otherwise} \end{cases}$$

We obviously have $h_i(\iota) < h_i(\kappa)$ whenever $\iota <_{\mathcal{O}_i} \kappa$. The height $h(\tau)$ of a non-progressive trace $\tau = (M_0, (\kappa_0, \kappa'_0)) \cdots (M_i, (\kappa_i, \kappa'_i)) \cdots$ is then defined by

$$h(\tau) = \sum_j (h_j(\kappa_j) - h_j(\kappa'_j))$$

$h(\tau)$ is finite, because τ is non-progressive. Now we define $d(i, \kappa) = \max(H(i, \kappa))$, where

$$H(i, \kappa) = \{h(\tau) \mid \tau \text{ is a trace following } \pi^i \text{ with } \tau(0) = (N_i, (\kappa_i, \kappa'_i)) \text{ and } \kappa_i = \kappa\}$$

In order to see that d is well-defined note that no non-progressive trace following π^i can have more than $m = n \cdot k^2$ progressing positions, where $n = |\mathcal{N}|$ is the number of nodes of \mathcal{P} and k is the number of distinct ordinal variables in \mathcal{P} . Any trace τ with more than m progressing positions must repeat some pair $\tau(j) =$

$\tau(k) = (N, (\kappa, \kappa'))$ with $j < k$, $N(\Gamma \vdash_{\mathcal{O}} \Delta)$ and $\kappa' <_{\mathcal{O}} \kappa$. This implies that $\tau' = (\tau(j)\tau(j+1)\cdots\tau(k-1))^\omega$ is a progressive trace following π^{i+j} , contradicting our assumption that no such trace exists. Hence, $H(i, \kappa)$ is bounded by $m \cdot l$, where $l = \max\{h_j(\kappa') \mid j \geq i \text{ and } \kappa' \in \mathcal{O}_j\}$, showing that d is well-defined.

Next, we show that d satisfies the following properties:

- (i) $d(i, \kappa) < d(i, \lambda)$ whenever $\kappa <_{\mathcal{O}_i} \lambda$,
- (ii) $d(i+1, \kappa) \leq d(i, \kappa)$ whenever $(N_i, N_{i+1}) \in \mathcal{E}$ and $\kappa \in |\mathcal{O}_i|$, and
- (iii) $d(i+1, \kappa) \leq d(i, \sigma(\kappa))$ whenever $(N_{i+1}, N_i, \sigma) \in \mathcal{R}$ and $\kappa \in |\mathcal{O}_{i+1}|$.

To see (i) assume that $\tau = (N_i, (\kappa, \kappa'))\tau'$ is a trace following π^i such that $h(\tau) = d(i, \kappa)$. Then $\tau'' = (N_i, (\lambda, \kappa'))\tau'$ is a trace following π^i with $h(\tau'') > h(\tau)$, hence $d(i, \lambda) \geq h(\tau'') > d(i, \kappa)$. For (ii) suppose $(N_i, N_{i+1}) \in \mathcal{E}$ is an edge of \mathcal{D} , $\kappa \in |\mathcal{O}_i|$ and $\tau = (N_{i+1}, (\kappa, \kappa'))\tau'$ is a trace following π^{i+1} such that $h(\tau) = d(i+1, \kappa)$. Then $\tau'' = (N_i, (\kappa, \kappa))\tau$ is a trace following π^i with $h(\tau'') = h(\tau)$, thus $d(i, \kappa) \geq h(\tau'') = d(i+1, \kappa)$. For (iii) suppose (N_{i+1}, N_i, σ) is a repeat in \mathcal{R} and that $\tau = (N_{i+1}, (\kappa, \kappa'))\tau'$ is a trace following π^{i+1} such that $h(\tau) = d(i+1, \kappa)$. Then $\tau'' = (N_i, (\sigma(\kappa), \sigma(\kappa)))\tau$ is a trace following π^i with $h(\tau'') = h(\tau)$, thus $d(i, \sigma(\kappa)) \geq h(\tau'') = d(i+1, \kappa)$.

We are now in a position to construct an infinite run $\Pi = (N_0, \rho_0) \cdots (N_i, \rho_i) \cdots$ following π . The construction will satisfy the invariants

- (J1) $\rho_i(\kappa) \geq d(i, \kappa)$ for all $\kappa \in |\mathcal{O}_i|$
- (J2) ρ_i respects \mathcal{O}_i

at each position $i \in \mathbb{N}$. We start by setting $\rho_0(\kappa) = d(0, \kappa)$ for each $\kappa \in |\mathcal{O}_0|$, which trivially satisfies (J1). ρ_0 also satisfies (J2) by (i) above.

Now suppose we have already constructed $(N_0, \rho_0) \cdots (N_i, \rho_i)$ such that (J1) and (J2) hold for i . We define ρ_{i+1} and show that it satisfies invariants (J1) and (J2). We distinguish two cases:

Case 1. (N_i, N_{i+1}) is an edge of \mathcal{D} . We proceed by a case analysis on the rule applied at N_i . Common to all cases is that we define $\rho_{i+1}(v) = \rho_i(v)$ for each $v \in \text{fv}(\mathcal{L}(N_i))$ (and, in particular, for $\kappa \in |\mathcal{O}_i|$). This implies

- (a) $\rho_{i+1}(\kappa) \geq d(i+1, \kappa)$ for all $\kappa \in |\mathcal{O}_i|$, and
- (b) $\rho_{i+1}(\iota) < \rho_{i+1}(\kappa)$ whenever $\iota <_{\mathcal{O}_i} \kappa$

by the induction hypothesis and (ii). Since we have $\mathcal{O}_i \subseteq \mathcal{O}_{i+1}$, this establishes (J1) and (J2) for all rules except $(\exists_{\mathcal{O}}\text{-L})$, $(\exists_{\mathcal{O}}^<\text{-L})$ and (OrdStr) . For the latter rules it remains to define ρ_{i+1} on any freshly introduced ordinal variables and to check invariants (J1) and (J2) for the additional cases concerning the fresh variables:

- $(\exists_{\mathcal{O}}\text{-L})$:** We set $\rho_{i+1}(\iota) = d(i+1, \iota)$, where ι is the fresh ordinal variable introduced by the rule. (J1) is satisfied by construction and (J2) is satisfied vacuously, since there are no cases involving ι .
- $(\exists_{\mathcal{O}}^<\text{-L})$:** We set $\rho_{i+1}(\iota) = d(i+1, \iota)$, where ι is the fresh ordinal variable introduced by the rule. (J1) is satisfied by construction. For (J2) let $\kappa \in |\mathcal{O}_i|$ such that $\iota <_{\mathcal{O}_{i+1}} \kappa$. By (i) we have $d(i+1, \iota) < d(i+1, \kappa)$ and by (a) $d(i+1, \kappa) \leq \rho_{i+1}(\kappa)$, so $\rho_{i+1}(\iota) < \rho_{i+1}(\kappa)$ as required.

(OrdStr): Let $m = \max\{\rho_i(\lambda) \mid \lambda \in |\mathcal{O}_i|\}$ and $\rho_{i+1}(\kappa) = d(i+1, \kappa) + m + 1$ for $\kappa \in |\mathcal{O}_{i+1}| - |\mathcal{O}_i|$. Invariant (J1) is clearly satisfied by construction. It remains to verify (J2). Suppose that $\iota <_{\mathcal{O}_{i+1}} \kappa$. Since the side condition of the rule guarantees that $\iota \in |\mathcal{O}_i|$ whenever $\iota <_{\mathcal{O}_{i+1}} \kappa$ and $\kappa \in |\mathcal{O}_i|$, there are two cases not already covered by (a) and (b). In the first case, where $\iota, \kappa \in |\mathcal{O}_{i+1}| - |\mathcal{O}_i|$, the result follows from (i). For the second case, where $\iota \in |\mathcal{O}_i|$ and $\kappa \in |\mathcal{O}_{i+1}| - |\mathcal{O}_i|$, we have $\rho_{i+1}(\iota) \leq m < \rho_{i+1}(\kappa)$ by the definitions of m and ρ_{i+1} .

Case 2. (N_{i+1}, N_i, σ) is a repeat in \mathcal{R} . We set $\rho_{i+1} = \rho_i \circ \sigma$. By induction hypothesis we have $\rho_i(\sigma(\kappa)) \geq d(i, \sigma(\kappa))$ and by (iii) $d(i, \sigma(\kappa)) \geq d(i+1, \kappa)$, hence (J1) holds. For (J2) suppose that $\iota <_{\mathcal{O}_{i+1}} \kappa$. Then $\sigma(\iota) <_{\mathcal{O}_i} \sigma(\kappa)$ by the definition of a repeat, thus we get $\rho_{i+1}(\iota) = \rho_i(\sigma(\iota)) < \rho_i(\sigma(\kappa)) = \rho_{i+1}(\kappa)$ from the induction hypothesis.

Continuing this construction ad infinitum yields an infinite run Π following π .

“ \Leftarrow ” For the opposite direction suppose there is a progressive trace

$$\tau = (N_i, (\kappa_i, \kappa'_i))(N_{i+1}, (\kappa_{i+1}, \kappa'_{i+1})) \cdots$$

following the suffix π^i of π . For a contradiction suppose further that there is an infinite run $\Pi = (N_0, \rho_0) \cdots (N_i, \rho_i) \cdots$ following π . Let $j \geq i$. Then we have $\rho_j(\kappa_j) \geq \rho_j(\kappa'_j)$, since ρ_j respects \mathcal{O}_j . We also have $\rho_j(\kappa'_j) = \rho_{j+1}(\kappa_{j+1})$ by the definition of a run, since $\kappa'_j = \kappa_{j+1}$ whenever $(N_j, N_{j+1}) \in \mathcal{E}$ is a tree edge and $\kappa'_j = \sigma(\kappa_{j+1})$ whenever $(N_{j+1}, N_j, \sigma) \in \mathcal{R}$ is a repeat of \mathcal{P} . Thus, there is an infinite chain $\rho_i(\kappa_i) \geq \rho_{i+1}(\kappa_{i+1}) \geq \cdots$ of ordinals, which strictly decreases at infinitely many positions, since τ is progressive. This contradicts the well-foundedness of the ordinals. \square

4.3. AUTOMATA-THEORETIC CHARACTERISATION

While the trace-based discharge condition is syntactic, it is – as it stands – still not suitable for practical application in a proof tool, since it is defined in terms of infinite objects. In order to obtain an implementable condition, we now turn to an automata-theoretic reformulation of the trace-based discharge condition.

Technically, this new condition will be realised using Büchi automata for which we introduce the following conventions. A Büchi automaton $\mathcal{A} = (A, Q, Q^0, \delta, F)$ is composed of an alphabet A , a finite set of states Q , a set of initial states $Q^0 \subseteq Q$, a transition relation $\delta \subseteq Q \times A \times Q$ and a set of accepting states $F \subseteq Q$. An infinite word $\sigma \in A^\omega$ is accepted by \mathcal{A} if there is a run $r \in Q^\omega$ over σ visiting F infinitely often. We denote by $L(\mathcal{A})$ the language accepted by \mathcal{A} . For more details we refer the reader to Thomas’ handbook chapter [15].

Essentially, the automata-theoretic characterisation looks for the existence of specific traces, which progress only at repeat nodes.

Definition 4.6. (Normal trace) A trace $\tau = (N_0, (\kappa_0, \kappa'_0)) \cdots (N_i, (\kappa_i, \kappa'_i)) \cdots$ is called *normal* if, for all i , node N_i is a repeat node whenever τ progresses at position i . \diamond




FIGURE 5. Example of a normal trace

Example 4.7. Figure 5 shows the the trace

$$\tau = (N_0, (\delta, \varepsilon))(N_1, (\alpha, \beta))(N_2, (\beta, \gamma))(N_3, (\gamma, \gamma))(N_4, (\kappa, \kappa)),$$

of the previous Example 4.3 using continuous lines. Replacing the continuous line with the dashed lines between positions 1 and 3 yields the normal variant

$$\tau' = (N_0, (\delta, \varepsilon))(N_1, (\alpha, \alpha))(N_2, (\alpha, \alpha))(N_3, (\alpha, \gamma))(N_4, (\kappa, \kappa)),$$

of τ , where progress of τ at positions 1 and 2 is deferred to the repeat node at position 3 in τ' .

Lemma 4.8. *Every trace τ can be transformed into a normal trace τ' such that τ is progressive if and only if τ' is progressive.*

Proof. (Sketch) Since $\mathcal{O}_i \subseteq \mathcal{O}_{i+1}$ whenever $(N_i, N_{i+1}) \in \mathcal{E}$, progress can be deferred to repeat nodes in the manner suggested by Example 4.7. As progress is only deferred to the next repeat but never lost, progressiveness is preserved by this transformation. \square

By Lemma 4.8 we may without loss of generality for condition (tDC) restrict our attention to the *normal* traces of \mathcal{P} . Based on this observation we construct two Büchi automata, \mathcal{B}_1 and \mathcal{B}_2 , over the alphabet \mathcal{R} of repeats.

The path automaton \mathcal{B}_1 : recognises those sequences of repeats that are traversed by paths of \mathcal{P} .

The progress automaton \mathcal{B}_2 : recognises sequences of repeats that are *potentially* connected through a normal trace; potentially, because this automaton tracks ordinal variable dependencies as in a normal trace, but completely ignores whether the sequence of repeats it accepts may be traversed by some path of \mathcal{P} .

The language inclusion $L(\mathcal{B}_1) \subseteq L(\mathcal{B}_2)$ then holds precisely if there is a normal trace along each infinite path of \mathcal{P} . Some auxiliary definitions prepare the formal definition of these two automata.

Definition 4.9. The relation $\rightarrow \subseteq \mathcal{R} \times \mathcal{R}$ on the set of repeats of \mathcal{P} is defined by $R \rightarrow R'$ if there is a path in the derivation tree \mathcal{D} of \mathcal{P} from the companion node of R to the repeat node of R' .

We also define $V_o = \bigcup \{|\mathcal{O}| \mid N(\Gamma \vdash_{\mathcal{O}} \Delta) \in \mathcal{N}\}$, the set of free ordinal variables of \mathcal{P} , and let r_π be the sequence of repeats traversed by a path π of \mathcal{P} .

Definition 4.10. The *path automaton* of \mathcal{P} is the Büchi automaton

$$\mathcal{B}_1 = (\mathcal{R}, Q_1, Q_1^0, \delta_1, F_1)$$

where $Q_1 = Q_1^0 = F_1 = \mathcal{R}$ and the transition relation $\delta_1 \subseteq Q_1 \times \mathcal{R} \times Q_1$ is defined by $\delta_1 = \{(R, R, R') \mid R \rightarrow R'\}$.

The following characterisation of the language accepted by \mathcal{B}_1 follows immediately from the definitions.

Lemma 4.11. $L(\mathcal{B}_1) = \{r_\pi \mid \pi \text{ an infinite path of } \mathcal{P}\}$.

Definition 4.12. The *progress automaton* of \mathcal{P} is the Büchi automaton

$$\mathcal{B}_2 = (\mathcal{R}, Q_2, Q_2^0, \delta_2, F_2)$$

where $Q_2 = Q_2^0 = (V_o \times \mathbf{2} \times V_o) \cup \{\diamond\}$, $F_2 = V_o \times \{1\} \times V_o$ and $\delta_2 \subseteq Q_2 \times \mathcal{R} \times Q_2$ is defined by $\delta_2 = \delta_2' \cup (\{\diamond\} \times \mathcal{R} \times Q_2)$ with

$$\delta_2' = \{((\iota, a, \kappa), (M, N, \sigma), (\kappa, b, \lambda)) \mid \sigma(\kappa) \leq_{\mathcal{O}_N} \iota \text{ and } a = 0 \Leftrightarrow \sigma(\kappa) = \iota\}$$

Note the presence of the state \diamond and the transitions from this state to any other state (including itself). Its role is to ensure that the language accepted by \mathcal{B}_2 is closed under prefixing with finite words over \mathcal{R} , reflecting the requirement in condition (tDC) that each infinite rooted path π has a trace following some *suffix* of π . Let us now illustrate these definitions with an example.




FIGURE 6. Example transition of automaton \mathcal{B}_2

Example 4.13. The upper part of Figure 6 shows a simplified version of the normal trace from Example 4.7. Since $\sigma_1(\alpha) <_{\mathcal{O}_0} \delta$ this trace gives rise to a transition $(\delta, 1, \alpha) \xrightarrow{R_1} (\alpha, j, \kappa)$ of automaton \mathcal{B}_2 for each $j \in \mathbf{2}$.

Definition 4.14. (aDC) A pre-proof \mathcal{P} satisfies condition (aDC) if

$$L(\mathcal{B}_1) \subseteq L(\mathcal{B}_2).$$

Theorem 4.15. *A pre-proof \mathcal{P} satisfies condition (tDC) if and only if \mathcal{P} satisfies condition (aDC). The latter condition can be checked in time $2^{\mathcal{O}(m^2 \log m)}$, where $m = n + r$ is the sum of the number of nodes n of \mathcal{P} and the number of ordinal variables r occurring free in the root sequent of \mathcal{P} .*

Proof. By Lemmas 4.8 and 4.11, it is sufficient to show for all infinite paths π that there is a progressive normal trace τ following a suffix of π if and only if there is an accepting run r of \mathcal{B}_2 on r_π . Accordingly, let $\pi = N_0 N_1 \cdots$ be an infinite path of \mathcal{P} with $N_i(\Gamma_i \vdash_{\mathcal{O}_i} \Delta_i)$.

“ \Rightarrow ” Consider a progressive normal trace

$$\tau = (N_k, (\kappa_k, \kappa'_k))(N_{k+1}, (\kappa_{k+1}, \kappa'_{k+1})) \cdots \quad (2)$$

following π^k for some $k \geq 0$. Let i_0, i_1, \dots be the positions where a repeat node appears on τ and let $R_j = (N_{i_j+1}, N_{i_j}, \sigma_j)$ for $j \geq 0$ be the corresponding repeats. We construct the infinite sequence

$$r = \diamond^p(\lambda_0, k_0, \lambda_1)(\lambda_1, k_1, \lambda_2) \cdots \quad (3)$$

where p is the number of repeat nodes appearing before position k on π and $\lambda_j = \kappa_{i_j}$ for $j \geq 0$. Since τ is a trace, we have $\lambda_j = \kappa_{i_j} \geq_{\mathcal{O}_{i_j}} \sigma_j(\kappa_{i_j+1})$. We also have $\kappa_{i_j+1} = \kappa_{i_{j+1}} = \lambda_{j+1}$, because τ is normal. We set $k_j = 0$ if $\sigma(\lambda_{j+1}) = \lambda_j$ and $k_j = 1$ otherwise. It is then not difficult to see that r is a run of \mathcal{B}_2 on r_π , which is accepting since τ is progressive.

“ \Leftarrow ” Suppose r is an accepting run of \mathcal{B}_2 on r_π of the form (3) above, let $r_\pi^p = R_0 R_1 \cdots$ and let i_j be the position of repeat $R_j = (N_{i_j+1}, N_{i_j}, \sigma_j)$ on π for each $j \geq 0$. We construct an infinite sequence τ of the shape 2 above by setting $k = i_0$ and

$$\begin{aligned} (\kappa_{i_j}, \kappa'_{i_j}) &= (\lambda_j, \sigma_j(\lambda_{j+1})) \\ (\kappa_l, \kappa'_l) &= (\lambda_{j+1}, \lambda_{j+1}) \quad \text{for } i_j + 1 \leq l \leq i_{j+1} - 1 \end{aligned}$$

for $j \geq 0$. By the definition of \mathcal{B}_2 we know that $\sigma_j(\lambda_{j+1}) \leq_{\mathcal{O}_{i_j}} \lambda_j$ for all $j \geq 0$. Because $\mathcal{O}_l \subseteq \mathcal{O}_{l+1}$ whenever (N_l, N_{l+1}) is a tree edge, it is then easy to see that τ is a normal trace following π^k , which is progressive since r is accepting.

It remains to justify the complexity claim. The standard way to check the inclusion $L(\mathcal{B}_1) \subseteq L(\mathcal{B}_2)$ is to complement \mathcal{B}_2 into $\overline{\mathcal{B}_2}$ and check the product $\mathcal{B}_1 \times \overline{\mathcal{B}_2}$ for emptiness. A pre-proof \mathcal{P} with $n = |\mathcal{N}|$ nodes can have at most n repeats. The number $|V_o|$ of ordinal variables in \mathcal{P} is bounded by $m = n + r$, where r is the number of free ordinal variables of the root sequent. This yields $\mathcal{O}(n)$ states for \mathcal{B}_1 and $\mathcal{O}(m^2)$ states for \mathcal{B}_2 . Complementing a Büchi automaton with n states can be done in time $2^{\mathcal{O}(n \log n)}$ [10]. Hence, the complementation of

\mathcal{B}_2 takes time $2^{\mathcal{O}(m^2 \log m)}$, which does not increase by computing the product with \mathcal{B}_1 and the subsequent linear time emptiness check. \square

5. RESTRICTED FORMS OF SYNTACTIC DISCHARGE

In this section we present two more restrictive syntactic discharge conditions and relate them to our new conditions as well as with those proposed in the literature. Let us consider an arbitrary but fixed pre-proof $\mathcal{P} = (\mathcal{D}, \mathcal{R})$.

5.1. DISCHARGE BASED ON STRONGLY CONNECTED SETS OF REPEATS

Definition 5.1. Let $R = (M, N, \sigma)$ be a repeat such that $M(\Gamma' \vdash_{\mathcal{O}'} \Delta')$ and $N(\Gamma \vdash_{\mathcal{O}} \Delta)$, and let $\kappa \in |\mathcal{O}'|$ be an ordinal variable. Then we say

- (1) R *preserves* κ if $\sigma(\kappa) \leq_{\mathcal{O}} \kappa$, and
- (2) R *progresses on* κ if $\sigma(\kappa) <_{\mathcal{O}} \kappa$.

A set of repeats $S \subseteq \mathcal{R}$ is called *strongly connected* if $(S, \rightarrow \cap (S \times S))$, the subgraph of $(\mathcal{R}, \rightarrow)$ induced by S , is strongly connected. Equivalently, one can say that there is a path π traversing exactly the repeats in S infinitely often.

Definition 5.2. (scDC) A pre-proof $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ satisfies *condition (scDC)* if for each strongly connected $S \subseteq \mathcal{R}$ there is an ordinal variable κ such that

- (1) some repeat $R \in S$ progresses on κ , and
- (2) each repeat $R' \in S$ preserves κ .

This condition is similar to the one described by Fredlund [6]. Schöpp and Simpson [12] use essentially the same condition as well, although their proof system is based on a different notion of approximation without ordinal variables.

Condition (scDC) can be reformulated automata-theoretically by replacing the trivial Büchi acceptance condition of the path automaton \mathcal{B}_1 of Definition 4.10 by a non-trivial Streett acceptance condition. A Streett automaton $\mathcal{A} = (\Sigma, Q, Q^0, \delta, \Omega)$ has the same components as Büchi automaton except that the acceptance condition is replaced by the *Streett acceptance condition* $\Omega = \{(L_i, U_i) \mid 1 \leq i \leq n\}$ consisting of a set of pairs of states. An infinite word σ is accepted by \mathcal{A} if there is a run $r \in Q^\omega$ such that, for all i , if r visits L_i infinitely often then it also visits U_i infinitely often. To capture condition (scDC) we define the Streett automaton $\mathcal{S} = (\Sigma, Q, Q^0, \delta, \Omega)$, where $\Sigma = Q = Q^0 = \mathcal{R}$ and $\delta = \{(R, R, R') \mid R \rightarrow R'\}$. The acceptance condition is $\Omega = \{(L_\kappa, U_\kappa) \mid \kappa \in V_o\}$, where V_o is the set of ordinal variables occurring free in \mathcal{P} and

$$\begin{aligned} L_\kappa &= \{R \in \mathcal{R} \mid R \text{ progresses on } \kappa\} \\ U_\kappa &= \{R \in \mathcal{R} \mid R \text{ does not preserve } \kappa\} \end{aligned}$$

Proposition 5.3. *A pre-proof \mathcal{P} satisfies condition (scDC) if and only if $L(\mathcal{S})$ is empty. The latter condition can be checked in time $\mathcal{O}(m^3)$, where $m = n + r$ is the sum of the number of nodes n of \mathcal{P} and the number of ordinal variables r occurring free in the root sequent of \mathcal{P} .*

Proof. The first part is not difficult to see from the definitions. The complexity of checking the emptiness of a Streett automaton is $\mathcal{O}((n+k)^2 \min(n,k))$, where n is the number of states and k is the number of accepting pairs [5]. The result follows, since in our case m is an upper bound of both the number of repeats and the number of ordinal variables in \mathcal{P} . \square

For a comparison of condition (scDC) with our previous condition (aDC), we define

$$\mathcal{B}_2^- = (\mathcal{R}, Q_2^-, Q_2^0 \cap Q_2^-, \delta_2 \cap Q_2^- \times \mathcal{R} \times Q_2^-, F_2 \cap Q_2^-)$$

to be the Büchi automaton obtained from $\mathcal{B}_2 = (\mathcal{R}, Q_2, Q_2^0, \delta_2, F_2)$ by restricting the sets of states and transitions to the set $Q_2^- = \{(\iota, j, \kappa) \in Q_2 \mid \iota = \kappa\}$. Condition (aDC-) then requires that $L(\mathcal{B}_1) \subseteq L(\mathcal{B}_2^-)$.

Proposition 5.4. *A pre-proof \mathcal{P} satisfies condition (scDC) if and only if it satisfies condition (aDC-).*

Proof. “ \Rightarrow ” Suppose \mathcal{P} satisfies (scDC) and let $r_1 = R_0 \cdots R_i \cdots$ be an accepting run of \mathcal{B}_1 on r_1 . We show that r_1 is accepted by \mathcal{B}_2 . Let S be the set of repeats occurring infinitely often in r_2 . Since S is strongly connected, there is an ordinal variable κ such that some $R \in S$ progresses on κ and all $R' \in S$ preserve κ . Since $\mathcal{G}(\mathcal{P})$ is finite there is a position k such that all R_j with $j \geq k$ belong to S . Define $r_2 = \diamond^k(\kappa, i_k, \kappa)(\kappa, i_{k+1}, \kappa) \cdots$, where, for each $j \geq k$, we set $i_j = 1$ if R_j progresses on κ and $i_j = 0$ otherwise. Then r_2 is a run of \mathcal{B}_2^- on r_1 , which is accepting, since there are infinitely many $j \geq k$ such that $R_j = R$. Thus, \mathcal{P} satisfies condition (aDC-).

“ \Leftarrow ” Suppose \mathcal{P} satisfies (aDC-) and let $S \subseteq \mathcal{R}$ be strongly connected. Then there is an accepting run $r_1 = R_0 R_1 \cdots$ of \mathcal{B}_1 on r_1 such that $\{R_i \mid i \geq 0\} = S$. By (aDC-) there is an accepting run $r_2 = (\kappa, i_0, \kappa)(\kappa, i_1, \kappa) \cdots$ of \mathcal{B}_2^- on r_1 , implying that condition (scDC) holds for S and ordinal variable κ . \square

Corollary 5.5. *If a pre-proof \mathcal{P} satisfies (scDC) then it satisfies condition (tDC).*

Proof. By Theorem 4.15, since $L(\mathcal{B}_2^-) \subseteq L(\mathcal{B}_2)$. \square

The following example shows that the converse of Corollary 5.5 does not hold in general.

Example 5.6. Let $\phi = \mu X(x). \exists z. X(z)$. The derivation in Figure 7 shows a pre-proof \mathcal{P} for the sequent $\phi(x), \phi(y) \vdash$. We write ϕ^κ for $\mu^\kappa X(x). \exists z. X(z)$. We have named the nodes for reference and omitted some intermediate nodes for a more compact presentation.

This pre-proof has one repeat $R = (N_2, N_4, \sigma)$ with $\sigma = [x'/y, \kappa'/\iota', \kappa/\iota, \iota'/\kappa]$. Recall that we identify formulas up to renaming of bound variables. There is a infinite normal trace τ^ω , where

$$\tau = (N_2, (\iota, \iota))(N_3, (\iota, \iota))(N_4, (\iota, \iota'))(N_2, (\kappa, \kappa))(N_3, (\kappa, \kappa))(N_4, (\kappa, \kappa)),$$

following the the suffix $(N_2 N_3 N_4)^\omega$ of the only infinite path of \mathcal{P} . The trace τ is progressive, since $\iota' <_{\mathcal{O}_4} \iota$ at node N_4 . Hence, \mathcal{P} satisfies condition (tDC)

$$\begin{array}{c}
\frac{N_0[\phi(x), \phi(y) \vdash]}{N_1[\phi'(x), \phi^\kappa(y) \vdash_{\iota, \kappa}]} \quad (\mu_1\text{-L}, \exists_O\text{-L}) \\
\frac{N_2[\exists x'. \phi'(x'), \phi^\kappa(y) \vdash_{\iota' < \iota, \kappa}]}{N_3[\exists x'. \phi'(x'), \exists y'. \phi^{\kappa'}(y') \vdash_{\iota' < \iota, \kappa' < \kappa}]} \quad (\mu^\kappa\text{-L}, \exists_O^\leq\text{-L}) \\
\frac{N_3[\exists x'. \phi'(x'), \exists y'. \phi^{\kappa'}(y') \vdash_{\iota' < \iota, \kappa' < \kappa}]}{N_4[\phi'(x'), \exists y'. \phi^{\kappa'}(y') \vdash_{\iota' < \iota, \kappa' < \kappa}]} \quad (\exists_I\text{-L})
\end{array}$$

FIGURE 7. Pre-proof distinguishing (tDC) from (scDC)

and is thus a proof for $\phi(x), \phi(y) \vdash$. On the other hand, repeat R does not preserve any ordinal variable according to Definition 5.1. Hence, \mathcal{P} fails to satisfy condition (scDC).

5.2. DISCHARGE USING INDUCTION ORDERS

We introduce an alternative discharge condition based on ordering the repeats of a pre-proof. Here, we restrict our attention to *simple* pre-proofs $\mathcal{P} = (\mathcal{D}, \mathcal{R})$, where for each repeat $(M, N, \sigma) \in \mathcal{R}$ there is a path from M to N in \mathcal{D} .

Definition 5.7. Let $R = (M, N, \sigma)$ and $R' = (M', N', \sigma')$ be two repeats in \mathcal{R} . The *structural dependency relation* $\leq_{\mathcal{P}}$ on repeats is defined by $R \leq_{\mathcal{P}} R'$ if the companion M of R lies on the path $\pi_{R'} = M' \cdots N'$ from the companion M' to the repeat node N' of R' . Let $\asymp_{\mathcal{P}} = \leq_{\mathcal{P}} \cup \leq_{\mathcal{P}}^{-1}$ be the symmetric closure of $\leq_{\mathcal{P}}$.

The following two lemmas establish some useful connections between the relations $\asymp_{\mathcal{P}}$, \rightarrow and strong connectedness.

Lemma 5.8. $R \asymp_{\mathcal{P}} R'$ if and only if $R \rightarrow R'$ and $R' \rightarrow R$ if and only if $\{R, R'\}$ is strongly connected.

Proof. Immediate from Definitions 4.9 and 5.7. \square

Lemma 5.9. Let $S \subseteq \mathcal{R}$ be strongly connected and let $R, R' \in S$. Then there is a $\asymp_{\mathcal{P}}$ -chain of repeats in S from R to R' , that is, there is a sequence $R_0 R_1 \cdots R_n$ of repeats in S such that $R = R_0$, $R' = R_n$ and $R_i \asymp_{\mathcal{P}} R_{i+1}$ for $0 \leq i < n$.

Proof. Suppose $R = (M, N, \sigma)$ and $R' = (M', N', \sigma')$ belong to the strongly connected set $S \subseteq \mathcal{R}$. It is sufficient to prove the conclusion under the additional assumption $R \rightarrow R'$. The general statement then follows by a routine induction. We first establish the following auxiliary property:

(P) if $R \rightarrow R'$ then either $R \asymp_{\mathcal{P}} R'$ or there exists $R'' \in S$ such that $R \rightarrow R''$, $R' \leq_{\mathcal{P}} R''$ and the companion M'' of R'' is a proper ancestor of M' in the derivation tree.

To see this, suppose that $R \rightarrow R'$, but not $R \asymp_{\mathcal{P}} R'$. Since \mathcal{P} is assumed to be simple, the companion M' lies on the path from M to N' , but not on the path

from M to N . As R and R' are in the same strongly connected set S , there is a path from R' to R in $(\mathcal{R}, \rightarrow)$, or equivalently, from M' back to M in \mathcal{P} . Hence, there must be some $R'' \in S$ such that $R' \rightarrow R''$ and whose companion node M'' lies above M' in the derivation tree. This implies that $R \rightarrow R''$ and $R' \leq_{\mathcal{P}} R''$.

Now suppose $R \rightarrow R'$. We show the existence of a $\succ_{\mathcal{P}}$ -chain from R to R' in S by induction on the length $l(R, R') = m$ of the path π from M to M' in \mathcal{P} . This is trivial for $m = 0$. For $m > 0$ we derive from property (P) that either $R \succ_{\mathcal{P}} R'$, in which case we are done, or there is some $R'' \in S$ such that $R \rightarrow R''$, $R'' \succ_{\mathcal{P}} R'$ and $l(R, R'') < m$. In the latter case, it follows from the induction hypothesis that there is a $\succ_{\mathcal{P}}$ -chain from R to R'' in S , which we complete into a $\succ_{\mathcal{P}}$ -chain from R to R' using $R'' \succ_{\mathcal{P}} R'$. \square

An induction order partially orders the repeats of a pre-proof. Repeats are required to be comparable under certain conditions.

Definition 5.10. (Induction Orders) A partial order (\mathcal{R}, \preceq) on the set of repeats is called an *induction order* for \mathcal{P} , if either $R \preceq R'$ or $R' \preceq R$ whenever

- (1) $R'' \preceq R$ and $R'' \preceq R'$ for some R'' (\preceq is *forest-like*), or
- (2) $R \succ_{\mathcal{P}} R'$ (\preceq *respects* $\succ_{\mathcal{P}}$)

A *labelled induction order* $(\mathcal{R}, \preceq, \delta)$ is an induction order (\mathcal{R}, \preceq) together with a map δ assigning an ordinal variable δ_R to each repeat $R \in \mathcal{R}$.

Under the mild restriction that each companion belongs to a unique repeat, the transitive closure of the structural dependency relation is an important special case of an induction order.

Proposition 5.11. *Let $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ be a pre-proof with unique companions, that is, no pair of distinct repeats share the same companion. Then the transitive closure of $\leq_{\mathcal{P}}$ is an induction order for \mathcal{P} .*

Proof. The relation $\leq_{\mathcal{P}}$ is reflexive and, by the uniqueness of companions, anti-symmetric. Hence, its transitive closure is a partial order. It is forest-like, since \mathcal{P} is a tree, and it respects $\succ_{\mathcal{P}}$, since it contains $\leq_{\mathcal{P}}$. \square

Definition 5.12. (ioDC) Let $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ be a simple pre-proof. We say that a labelled induction order $(\mathcal{R}, \preceq, \delta)$ *discharges* \mathcal{P} if for all $R \in \mathcal{R}$

- (1) R progresses on δ_R , and
- (2) R preserves $\delta_{R'}$ whenever $R \preceq R'$.

Pre-proof \mathcal{P} satisfies *condition (ioDC)* if there is a labelled induction order discharging \mathcal{P} .

Since any partial order that linearly orders the repeats of each strongly connected component of \mathcal{P} is an induction order, condition (ioDC) subsumes the original discharge condition (DC) proposed by Dam and Gurov [4]. For forest-like induction orders condition (ioDC) is equivalent to the condition given by Schöpp [11] as the following lemma will show. However, by relying on the structural dependency relation our new definition of induction order is more local in the sense that it avoids the quantification over all strongly connected subsets of

repeats of a pre-proof. This makes it easier to check whether a given partial order on the set of repeats is an induction order.

Lemma 5.13. *A forest-like partial order (\mathcal{R}, \preceq) is an induction order if and only if each strongly connected $S \subseteq \mathcal{R}$ has a \preceq -greatest element.*

Proof. “ \Rightarrow ” Suppose that (\mathcal{R}, \preceq) is an induction order. Let $S \subseteq \mathcal{R}$ be strongly connected. In order to see that S has a \preceq -greatest element, it is sufficient to show that any two $R, R' \in S$ have an upper bound in S , that is, $R \preceq \hat{R}$ and $R' \preceq \hat{R}$ for some $\hat{R} \in S$. Suppose $R, R' \in S$. By Lemma 5.9 there is a sequence $R_0 R_1 \cdots R_n$ of repeats in S such that $R = R_0$, $R' = R_n$ and $R_i \succ_{\mathcal{P}} R_{i+1}$ for $0 \leq i < n$. As (\mathcal{R}, \preceq) respects $\succ_{\mathcal{P}}$ we also have $R_i \preceq R_{i+1}$ or $R_{i+1} \preceq R_i$ for $0 \leq i < n$. Using the fact that (\mathcal{R}, \preceq) is forest-like a routine induction on n shows that there is an upper bound \hat{R} of R and R' in S .

“ \Leftarrow ” Suppose that each strongly connected $S \subseteq \mathcal{R}$ has a \preceq -greatest element and let R and R' be two repeats with $R \succ_{\mathcal{P}} R'$. Then $R \preceq R'$ or $R' \preceq R$ as required, since $S = \{R, R'\}$ is strongly connected by Lemma 5.8. \square

The next result shows that for simple pre-proofs discharge based on induction orders is equivalent to discharge based on strongly connected sets of repeats.

Theorem 5.14. *Let $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ be a simple pre-proof. Then \mathcal{P} satisfies condition (ioDC) if and only if it satisfies condition (scDC).*

Proof. “ \Rightarrow ” Suppose $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ satisfies (ioDC) witnessing the labelled induction order $(\mathcal{R}, \preceq, \delta)$ and let $S \subseteq \mathcal{R}$ be strongly connected. Then S has a \preceq -greatest element R by Lemma 5.13. By the definition of discharge R progresses on δ_R and all $R' \in S$ preserve δ_R . Hence, \mathcal{P} satisfies condition (scDC).

“ \Leftarrow ” Suppose $\mathcal{P} = (\mathcal{D}, \mathcal{R})$ satisfies (scDC). We iteratively construct a labelled induction order $(\mathcal{R}, \preceq, \delta)$ as follows. We start with the set \mathcal{S}_0 partitioning \mathcal{R} into its strongly connected components. At step i we pick $S_i \in \mathcal{S}_i$ and then an $R_i \in S_i$ such that R_i progresses on some ordinal variable κ_i and all $R \in S_i$ preserve κ_i . Since \mathcal{P} satisfies (scDC), such R_i and κ_i exist. Then we set $\mathcal{S}_{i+1} = (\mathcal{S}_i - \{S_i\}) \cup D$, where D is obtained as the partitioning of $S_i - \{R_i\}$ into its strongly connected components. This process terminates after $n = |\mathcal{R}|$ iterations, since at each step i repeat R_i is removed from $\bigcup \mathcal{S}_i$.

We define δ by $\delta(R_i) = \kappa_i$ and $R_i \preceq R_j$ if $S_i \subseteq S_j$. By the hierarchical nature of the construction (\mathcal{R}, \preceq) is certainly a forest-like partial order and, moreover, for each strongly connected $S \subseteq \mathcal{R}$ there is a unique S_k such that $R_k \in S \subseteq S_k$. This implies that $S_i \subseteq S_k$ for any $R_i \in S$ and thus R_k is \preceq -greatest in S . Hence, (\mathcal{R}, \preceq) is an induction order for \mathcal{P} by Lemma 5.13. Note that any repeat R progresses on δ_R by construction. Also, for any R_i with $R_i \preceq R_j$, we have $R_i \in S_i \subseteq S_j$, so R_i preserves δ_{R_j} . Thus, $(\mathcal{R}, \preceq, \delta)$ discharges \mathcal{P} . \square

6. CONCLUSIONS

We have studied a Gentzen-style proof system for the μ -calculus which is based on circular proofs. In particular, we have investigated several discharge conditions which externally justify the well-foundedness of inductive reasoning embodied in these proofs. Starting from the natural semantic condition proposed by Dam and Gurov [4], we have, based on the notion of traces, given a syntactical condition which characterises the semantic one for any given pre-proof. While this condition is purely syntactical, it is still not suitable for implementation as its definition directly refers to infinite objects. Therefore, we have also elaborated an algorithmic formulation in terms of a language inclusion between two Büchi automata.

Next, for a detailed comparison with previously known discharge conditions, we have focused our attention on two simpler discharge criteria. In particular, we have considered two levels of restrictions with respect to our general condition:

- (1) Restrict to normal traces that track the behaviour of a single ordinal variable, disallowing its renaming at repeats; this leads to condition (scDC), similar to those in [6, 12], requiring that we find, for each strongly connected subgraph, an ordinal variable and a repeat that progresses on this variable while the other repeats preserve it. This condition can be formulated as an emptiness problem of a Streett automaton.
- (2) Additionally restrict the form of pre-proofs to simple ones, where each repeat node is reachable in the proof tree from its companion node; this allowed us to organise the repeats of a pre-proof into a partial order, called induction order, and formulate a new condition, called (ioDC), which imposes progress and preservation conditions on each repeat according to its position in the induction order. This condition is close to those in [4, 11], but avoids a quantification over strongly connected subsets of repeats.

Our comparison showed that condition (scDC) and (ioDC) are equivalent for simple pre-proofs and condition (scDC) is in general strictly stronger than our trace-based condition on a *fixed* pre-proof. However, an important open question concerns the proof-theoretical strength of the different conditions. More precisely, it is currently unclear to us whether there are sequents that can be proved using the trace-based condition, but for which no proof exists if we restrict ourselves to using a simpler discharge criterion. We are inclined to think that this is not the case. But, while it might not be too difficult to show that any pre-proof can be unfolded into a simple one (with a potentially exponential blow-up due to the loss of sharing), the proof that we can dispense with renaming of ordinal variables at repeats seems more involved.

We would like to add some remarks regarding the practical application of our results. First, the exponential complexity of the general automata-based discharge condition (aDC) seems to discourage its use in favour of the more tractable condition (scDC) based on strongly connected components. We do not currently know whether there is a polynomial algorithm for the general condition, a question that

is left for future work. Second, the general condition is weaker and thus qualifies more pre-proofs as proofs, which can be an advantage in automatic proof search. However, it is unclear whether this difference frequently shows up in practise. Some experimentation is needed in order to clarify these issues. Third, in a tool implementation it is desirable to check discharge conditions incrementally in order to detect failure to discharge as soon as possible. Although the automata-based conditions (aDC) and (aDC-) can be used to this effect on the partially constructed proof structure, the need to complement the second automaton each time a new repeat is added does not support incremental checking very well. The reformulation of condition (scDC) as an emptiness problem of a Streett automaton is certainly easier to adapt for incremental verification.

Finally, it is important to observe that the dependency of our results on the μ -calculus itself is very limited. The μ -calculus was chosen here as a suitable minimal context in which to study global induction, but all that our induction mechanisms rely on is that the object language includes a form of inductive definition, which can be augmented by a corresponding notion of approximation. Given these ingredients it should, in principle, be possible to turn almost any deductive system using local induction rules into one based on global induction by replacing the local induction rules by appropriate versions of fixed point and ordinal rules. Some examples that merit a closer inspection are the inclusion of regular languages, inductive type theories and Hoare logics for recursive procedures. In a related paper [13], we investigate the relation between a proof system based on a local well-founded induction rule on the ordinals and one based on global induction as studied here. We establish their equivalence by giving proof translations in each direction.

We would like to thank Dilian Gurov, Lars-åke Fredlund and Alex Simpson for interesting and fruitful discussions on the subject of this paper. We are also grateful to the two anonymous referees for their very detailed and constructive comments.

REFERENCES

- [1] T. Arts, M. Dam, L. Fredlund, and D. Gurov. System description: Verification of distributed Erlang programs. In *Proc. CADE'98*, volume 1421 of *Lecture Notes in Artificial Intelligence*, pages 38–41, 1998.
- [2] J. Bradfield and C. Stirling. Local model checking for infinite state spaces. *Theoretical Computer Science*, 96:157–174, 1992.
- [3] M. Dam. Proving properties of dynamic process networks. *Information and Computation*, 140:95–114, 1998.
- [4] M. Dam and D. Gurov. μ -calculus with explicit points and approximations. *Journal of Logic and Computation*, 12(2):43–57, 2002. Previously appeared in Fixed Points in Computer Science, FICS 2000.
- [5] E. A. Emerson and C. L. Lei. Modalities for model checking: branching time strikes back. *Science of Computer Programming*, 8:275–306, 1987.
- [6] L. Fredlund. *A Framework for Reasoning about Erlang Code*. PhD thesis, Royal Institute of Technology, Stockholm, Sweden, 2001.

- [7] D. Kozen. Results on the propositional μ -calculus. *Theoretical Computer Science*, 27:333–354, 1983.
- [8] D. Niwiński and I. Walukiewicz. Games for the μ -calculus. *Theoretical Computer Science*, 163:99–116, 1997.
- [9] D. Park. Finiteness is mu-ineffable. *Theoretical Computer Science*, 3(2):173–181, 1976.
- [10] S. Safra. On the complexity of ω -automata. In *29th IEEE Symposium on Foundations of Computer Science*, pages 319–327, 1988.
- [11] U. Schöpp. Formal verification of processes. Master’s thesis, University of Edinburgh, 2001.
- [12] U. Schöpp and A. Simpson. Verifying temporal properties using explicit approximants: Completeness for context-free processes. In *Foundations of Software Science and Computational Structures (FoSSaCS 02), Grenoble, France*, volume 2303 of *Lecture Notes in Computer Science*, pages 372–386. Springer, 2002.
- [13] C. Sprenger and M. Dam. On the structure of inductive reasoning: Circular and tree-shaped proofs in the μ -calculus. In A. Gordon, editor, *Foundations of Software Science and Computational Structures (FoSSaCS 03), Warsaw, Poland, April 7-11, 2003*, volume 2620 of *Lecture Notes in Computer Science*, pages 425–440. Springer, 2003.
- [14] C. Stirling and D. Walker. Local model checking in the modal μ -calculus. *Theoretical Computer Science*, 89:161–177, 1991.
- [15] W. Thomas. Automata on infinite objects. In J. van Leeuwen, editor, *Handbook of Theoretical Computer Science*, volume B, pages 133–191. Elsevier Science Publishers, Amsterdam, 1990.

Communicated by (The editor will be set by the publisher).
(The dates will be set by the publisher).